

Class 2 Lab*

Sequence Alignment & Database Searching (Pt. 1)

Barry Grant

2025-12-16

i Instructions

Save this document to your computer and open it in a PDF viewer such as Preview (available on every mac) or Adobe Acrobat Reader ([free for PC and Linux](#)). Be sure to add your name and UC San Diego personal identification number (PID) and email below before answering all questions in the space provided.

Student Name

Patrick Nguyen

UCSD PID

A17680785

UCSD Email

pdn004@ucsd.edu

Learning Objectives

By the end of this lab, you will be able to:

1. **Explain** how dot plot parameters (window size, step size, and match stringency) affect the visualization of sequence similarity and the trade-off between sensitivity and specificity.
2. **Perform** pairwise sequence alignment by hand using the Needleman-Wunsch dynamic programming algorithm with specified scoring parameters.
3. **Interpret** the output of a completed scoring matrix to identify optimal alignments and recognize when multiple optimal solutions exist.
4. **Use** NCBI BLAST to search protein databases for homologous sequences, applying appropriate filters and database selections to refine results.
5. **Evaluate** BLAST results critically, considering E-values, percent identity, and query coverage to assess the biological significance of reported matches.

*<http://thegrantlab.org/teaching/>

Overview:

Aligning novel sequences with previously characterized genes or proteins provides important insights into their common attributes and evolutionary origins.

In sections 1, 2 and 3 of this hands-on session we will first explore the principles and methods underlying the computational comparison and alignment of biomolecular sequences.

In section 4 we explore how these methods are used to search databases to identify homologous sequences (i.e. finding evolutionary related genes or proteins that are descended from a common ancestor).

In section 5 we highlight the detection limits of conventional BLAST. This sets the scene for introducing more sensitive (but often more time consuming) approaches including Profiles, PSI-BLAST and Hidden Markov Models (HMMs).

Section 1: Dot Plot Parameters

Dot plots are a simple graphical approach for the visual comparison of two sequences. They have a long history (see [Maizel and Lenk 1981](#) and references therein) and entail placing one sequence on the vertical axis of a 2D grid (or matrix) and the other on the horizontal.

In its simplest form, a dot is placed where the horizontal and vertical sequence values match. More elaborate forms use 'sliding windows' composed of multiple characters and a threshold value, or 'match stringency' for two windows to be considered as matched.

Visit our very own simple dot plot web-app ([Link1](#) or it's mirror [Link2](#)) and get a feel for how altering these major dot plot parameters change the displayed protein and DNA dot plots.

N.B. Note the questions listed on the web page (also found below) and add your answers in the space provided on the next page.

BGGN-213: Dot Plot Comparison of Two Sequences

Dot plots are a simple graphical approach for the visual comparison of two sequences. They have a long history (see [Meisel and Lewis 1951](#)) and references therein) and entail placing one sequence on the vertical axis of a 2D grid (or matrix) and the other on the horizontal. In its simplest form, a dot is placed where the horizontal and vertical sequence values match. That is a dot is produced at position (i, j) if character number i in the first sequence is the same as character number j in the second sequence. More elaborate forms use 'sliding windows' composed of multiple characters and a threshold value, or 'match stringency' for two windows to be considered as matched.




Figure 1: Interactive dot plot web app to explore fundamental concepts of biomolecular sequence comparison

Q1 Why does the DNA sequence have more dots than the protein sequence plot?

Tip

What do you know about DNA composition vs protein composition?

Protein has a larger ladder (20 possible nucleotides) than DNA (4 possible nucleotides)

Q2 How can we increase the signal to noise ratio?

Tip

Signal in this case means correct matches that we actually want to highlight and noise means spurious matches that we don't want.

signal to noise ratio can be increased by increasing the window size and match stringency proportionally.

Q3 What does a 'Match stringency' larger than 'Window size' yield and why?

a "match stringency" larger than window size results in dot plot not being produced. A window size smaller than a match stringency cannot show the amount of matches asked by stringency larger than the window.

Q4 What are the major weaknesses of this approach?

Tip

Is your inner nerd happy with this approach? How would you use it to determine if a second set of sequences was more similar to each other than a first set of sequences?

Dot plots are not very quantitative and are only mostly qualitative.

Section 2: Needleman-Wunsch Alignment

Sequence alignment methods often use something called a 'dynamic programming' algorithm that can be usefully considered as an extension of the dot plot approach. Here we have two sample sequences, and we'd like to use the Needleman-Wunsch algorithm discussed in class to align them. Feel free to use the classroom white-boards and/or pen and paper and attach a photo to this PDF for gradescope.

Q5 Using a **match score** of +2, a **mismatch score** of -1, and a **gap score** of -2. Fill in the table below (or use pen and paper) for the following two sequences:

Sequence 1: ATTGC

Sequence 2: AGTTC

From your completed table (see template below) what is the **optimal score** and corresponding **alignment** (with one sequence above the other)?

Tip

It can be hard to store the all important progress arrows in the PDF version of this document and thus you may prefer to use your own paper (or white-board) version that you can take a photo off for upload to gradescope.

Optimal Score:

4

Alignments:

Alignment 1:

Sequence 1=A - T T G C A

Sequence 2=A G T T - C T

Alignment 2:

Sequence 1=A T T G C

Sequence 2=A G T T C

		A	G	T	T	C
	0					
A						
T						
T						
G						
C						

Section 3: Practice makes perfect

Again use the Needleman-Wunsch algorithm discussed in class to align the following sequences:

Q6 Using a **match score** of +2, a **mismatch score** of -1, and a **gap score** of -2. Fill in the table below and translate it into a alignment. What is the optimal score for this alignment? **Is there one unique alignment with this score?**


Sequence 1: TATAG
Sequence 2: GTTAC

Optimal Score:

1

Alignments:

Alignment 1:
Sequence 1=T A T A G
Sequence 2=G T T A C
Alignment 2:
Sequence 1= - T A T G G
Sequence 2= G T - T A C



		G	T	T	A	C
	0					
T						
A						
T						
A						
G						

Section 4: Finding homologous sequence

Your collaborators found a protein while working on a fly species and have asked you to see if there are any human homologs.

```
>fly_protein
MDNHSSVPWASAASVTCLSLDAKCHSSSSSSSSSSKAASSISAIPQEETQTMRHIAHTQRCLSRSLTSLVAL
LLIVLPMVFPSPAHSCGPGRGLGRHRARNLYPLVLKQTIPNLSEYTNASAGPLEGVIRRDSPKFKDLVPNY
NRDILFRDEEGTGADRLMSKRCCKEKLNVLAYSVNNEWPGIRLLVTESWDEDYHHGQESLHYEGRAVTIAT
SDRDQSKYGMRLARLAVEAGFDWVSYVSRRIYCSVKSDSSISSHVHGCFTPESTALLESQVRRKPLGELSI
GDRVLSMTANGQAVYSEVILFMDRNLEQMNFVQLHTDGGAVLTVTPAHLVSVWQPESQKLTFFVFADRIE
EKNQVLVRDVETGELRPQRVVKVGSVRSKGVVAPLTREGTIVVNSVAASCYAVINSQSLAHWGLAPMRL
STLEAWLPAKEQLHSSPKVVSSAQQQNGIHVYANALYKVKDYVLPQSWRHD
```

Q7 Using the default settings for NCBI BLAST, can you find any homologs for this protein in Humans?

Tip

Try using the LIMITS and FILTERING options we covered in the last lab.

no homologs found

Q8 Try changing the database to *refseq_protein*. From the results, select a few proteins and find the common name for the species. What trend do you notice as you move down the results list?

Tip

Search google for the species name and use the taxonomy tab on your NCBI BLAST results page.

hedgehog protein

Q9 Finally, try also limiting the search to only *H. Sapiens*. What function do these proteins have?

Tip

You can simply type the *Taxon ID 9606* in the "Organism" box.

calcium ion binding, protein binding, pepitase activity, patched binding, zinc ion binding

Q10 What function do you think this protein performs for your collaborators' organism?
guiding embryonic development, affecting patterning tissues

OPTIONAL EXTENSION

We will revisit this problem and introduce approaches with greater sensitivity (i.e. ability to find more remote homologues) in the next lab.

Section 5: The limits of using BLAST for remote homologue detection

Let's return to the HBB protein that we explored in a previous class and see if we can find distantly related myoglobin and neuroglobin using HBB as a BLAST query.

```
>gi|4504349|ref|NP_000509.1| hemoglobin subunit beta [Homo sapiens]
MVHLTPEEKSAVTALWGKVVNDEVGGEALGRLLVVYPWTQRFFESFGDLSTPDAVMGNPKVKAHGKKVLG
AFSDGLAHLDNAKGTFAITLSELHCDKLHVDPENFRLLGNVLCVLAHFGKEFTPPVQAAYQKVVAGVAN
ALAHKYH
```

After selecting **blastp** and entering the sequence, be sure to change the search database to **"refseq-protein"** and restrict our search organism to only **humans** (taxid: 9606). This will help focus our results to highlight distant homologs in humans.

Q11 What homologs did you find with this simple blastp search? Note their percent identities, coverage and E-values.

Now we could try changing the **Algorithm parameters** on the submission page to increase the number of hits reported. To do this you can click on the **Edit and Resubmit** link at the top left of your results page.

Q12 Try increasing the Expect threshold for your blasts search (e.g. to 2000). What new hits were reported? What about their alignment statistics? Do you trust these matches? Did you find myoglobin?



Q13: What one part of this exercise or associated lecture material is still confusing? If appropriate please also indicate the question number from this document and answer the question in the following anonymous form: [Muddy_Point_Assesment_Form](#) Your comments will let us know which material needs to be further clarified and will help us gain stronger control of the material in this course. Thank you!

Discussion

Many useful 'rules of thumb' are expressed in terms of percent identity. If two proteins have more than 45% identical residues in their optimal alignment they typically have very similar structures and are likely to have a similar function. If two proteins have more than 25% identical residues (but less than 45% identity), they are likely to have a similar general folding pattern. Note that we will expand on the basis of this important *sequence > structure > function* relationship in the next lab where we also explore the detection limits of standard BLAST searches for identifying remote homologs.

Observations of a lower degree of sequence similarity cannot however rule out homology. Our very own late [Russ Doolittle](#) defined the region between 18-25% sequence identity as the “twilight zone” in which the suggestion of homology is tantalizing but dangerous. Below the **twilight zone** is a region where pairwise sequence alignments tell us very little - sometimes called the “midnight zone”.

Our next class will introduce more advanced topics including profile and structure based approaches that can delve deeper into these important, but often hard to detect, sequence-structure-function relationships.